

## ANALYSIS OF A MULTIPHASE FREE BOUNDARY PROBLEM

Ahlem Abdelouahab and Sabri Bensid

*Communicated by J.I. Díaz*

**Abstract.** In this paper, we investigate a free boundary problem relevant in several applications, such as tumor growth models. Our problem is expressed as an elliptic equation involving discontinuous nonlinearities in a specified domain with a moving boundary. We establish the existence and uniqueness of solutions and provide a qualitative analysis of the free boundaries generated by the nonlinear term (inner boundaries). Furthermore, we analyze the dynamics of the outer region boundary. The final result demonstrates that under certain conditions, our problem is solvable in the neighborhood of a radial solution.

**Keywords:** discontinuous nonlinearity, free boundary, perturbation, tumor growth.

**Mathematics Subject Classification:** 34R35, 35J25, 92B05, 35R35.

### 1. INTRODUCTION

In various contexts, free boundaries often take the form of curves or surfaces whose exact positions are initially undetermined, serving as interfaces between regions with differing properties. An example is found in tumor growth models, where the boundary of the tumor region (separating it from healthy tissue) is not given a priori and changes during the process. This lack of knowledge requires a major study of the free boundary.

Over the last decades, several mathematical models have been proposed to describe the growth of tumor spheroids. The starting point in the study of free boundary problems arising in tumor growth models is due to the pioneering papers of Greenspan [20, 21]. Later, with a rigorous analysis, Byrne and Chaplain [8, 9] extended similar results to the case of discontinuous nonlinearity and recently, Friedman and collaborators in a series of papers [12, 15, 16, 18] develop mathematical techniques to analyze the existence, uniqueness and bifurcation of solutions. For the historical development of mathematical cancer modeling, we refer the reader to [2].

Inspired by the previous relevant works, the authors investigate in [1], for  $\varepsilon > 0$  the following problem:

$$\begin{cases} \Delta u = \lambda(\varepsilon + (1 - \varepsilon)H(u - \mu)) & \text{in } \Omega(t), \\ u = \bar{u}_\infty & \text{on } \partial\Omega(t), \end{cases} \quad (1.1)$$

where  $\mu$  is the critical value for which the tumor  $\Omega(t)$  developed a region of slow growth in its center due to lack of nutrients. Note that the problem (1.1) appears in other contexts like Budyko climate models. We refer the reader to [4] and [5] for more details.

Spherical coordinates, which are well suited to represent the spherical configuration of tumors are useful in analyzing our problem. So, let  $u := u(r, t)$  be the concentration nutrient in  $\Omega(t)$  where  $r = |x|$ . The dynamic of the outer tumor radius denoted by  $R(t)$  is governed by

$$\begin{aligned} \frac{d}{dt} \left( \frac{4\pi R^3(t)}{3} \right) &= \int \int \int_{\Omega(t)} S(u)r^2 \sin \theta \, d\theta \, d\phi \, dr \\ &\quad - \int \int \int_{\Omega(t)} N(u)r^2 \sin \theta \, d\theta \, d\phi \, dr, \end{aligned} \tag{1.2}$$

where  $S(u)$  and  $N(u)$  are the proliferation and the mortality rate functions.

The purpose of this work is to investigate the following problem:

$$\begin{cases} \Delta u = \lambda \left( \varepsilon H(u - \mu_1) + (1 - \varepsilon) H(u - \mu_2) \right) & \text{in } \Omega(t), \\ u = \bar{u}_\infty & \text{on } \partial\Omega(t), \end{cases} \tag{1.3}$$

where  $\Omega(t) \subset \mathbb{R}^3$  is the domain at time  $t > 0$  with a moving boundary  $\partial\Omega(t)$ ,  $\varepsilon \in (0, 1)$ ,  $\bar{u}_\infty, \lambda, \mu_1, \mu_2$  are positive parameters such that  $\mu_1 < \mu_2 < \bar{u}_\infty$  and  $H$  is the Heaviside function, i.e.

$$H(t) = \begin{cases} 1, & \text{for } t \geq 0, \\ 0, & \text{for } t < 0. \end{cases}$$

So, the present paper can be considered as a natural continuation of the previous paper by the authors [1] in which the same program of research was devoted to the one discontinuous nonlinearity. Here, we consider the sum of discontinuous nonlinearities, and as we shall see, our discontinuities generate free boundaries which are not given explicitly like [1] and this requires more attention. The challenge is to develop mathematical concepts that can handle the complexity of the problem.

In this work, we focus the study on the following multiphase free boundary problem

$$\begin{cases} \frac{1}{r^2} \frac{\partial}{\partial r} \left( r^2 \frac{\partial u}{\partial r} \right) = \lambda \left( \varepsilon H(u - \mu_1) + (1 - \varepsilon) H(u - \mu_2) \right), & 0 < r < R(t), t > 0, \\ u(R(t), t) = u_\infty, \frac{\partial u}{\partial r}(0, t) = 0 & \text{for } t > 0, \\ R^2(t) \frac{dR(t)}{dt} = \int_0^{R(t)} S(u)r^2 dr - \int_0^{R(t)} N(u)r^2 dr, \\ R(0) = R_0, \end{cases} \tag{1.4}$$

where  $\mu_1, \mu_2$  are the critical values for which tumor goes from one phase to another verifying  $\mu_1 < \mu_2 < u_\infty$  and

$$S(u) = \lambda(\varepsilon H(u - \mu_1) + (1 - \varepsilon) H(u - \mu_2)), \quad N(u) = \eta_1 H(u - \mu_1) + \eta_2 H(\mu_1 - u),$$

where  $\eta_1, \eta_2$  are positive constants.

Let us introduce the notations:

$$\omega_{\mu_i}(t) = \{u(r, t) < \mu_i\}, \quad \omega_{\mu_i} = \{u(r) < \mu_i\}$$

and

$$\Gamma_{\mu_i}(t) = \partial\omega_{\mu_i}(t), \quad \Gamma_{\mu_i} = \partial\omega_{\mu_i},$$

where  $\mu_i \in (0, \bar{u}_\infty)$  for  $i = 1, 2$ .

Hence, the discontinuous nonlinearities generate two free boundaries namely  $\Gamma_{\mu_1}(t)$  and  $\Gamma_{\mu_2}(t)$  in addition to the outer boundary  $\partial\Omega(t)$  (moving boundary). This is the first time that the problem (1.4) is considered. One of the main difficulties in problem (1.4) is that the free boundaries are not explicitly given (contrary to the paper [1]).

First, for the stationary solution and regarding the position of  $u$  with respect to  $\mu_1$  and  $\mu_2$  dealing with the three phases, we give the existence of a stationary solution for problem (1.4). Moreover, we show the existence of their corresponding free boundaries given by

$$u(r_i) = \mu_i, \quad i = 1, 2.$$

We further obtain the long time behavior of all transient solutions with the form  $(u(r, t), R(t))$ . Finally, we will assume that  $\Omega(t)$  is a perturbation of the ball  $B(0, R(t))$ . More precisely, we assume what follows:

$$\begin{aligned} \text{The boundary } \partial\Omega(t) \text{ can be parameterized as } R(t) + \beta(\theta) \text{ for } t > 0, \\ \text{where } \beta \in C^2(S), \theta \in S \text{ and } S \text{ is the unit sphere.} \end{aligned} \tag{C}$$

Using this parametrization of our free boundaries which are the unknowns of our problem, we reduce the study to the solvability of a nonlinear integral equation and show the existence of functions namely  $b_i$  such that

$$u(r_i(t) + b_i(\theta), \theta) = \mu_i, \quad i = 1, 2, t > 0.$$

Moreover, this perturbation method gives the existence of exceptional free boundaries in the case of a stationary solution with the form

$$u(r_i(t) + h(\theta), \theta) = \mu_i, \quad i = 1, 2,$$

where  $h \in C(S)$ . This result is far from obvious.

The structure of the rest of this paper is arranged as follows. In Section 2, we study the stationary solutions when  $\Omega = B(0, R(t))$  with different phases. Section 3 is devoted to establishing the existence of a transient solution and studying its asymptotic behavior. In Section 4, using perturbation, we reformulate our free boundary problem and prove the existence of free boundaries through the use of local methods. Finally, some comments are given.

2. STATIONARY SOLUTIONS

This section deals with the existence of stationary solutions for the problem (1.4). Hence, we consider the following problem:

$$\begin{cases} \frac{1}{r^2} \frac{\partial}{\partial r} \left( r^2 \frac{\partial u}{\partial r} \right) = \lambda \left( \varepsilon H(u - \mu_1) + (1 - \varepsilon) H(u - \mu_2) \right), & 0 < r < R, \\ u(R) = u_\infty, \frac{\partial u}{\partial r}(0) = 0, \end{cases} \tag{2.1}$$

and the integral equation

$$\frac{1}{R^2} \left( \int_0^R S(u) r^2 dr - \int_0^R N(u) r^2 dr \right) = 0. \tag{2.2}$$

We recall that the nonnegative solution of (2.1) is strictly convex function such that

$$\min_{0 < r < R} u(r) = u(0).$$

The sets  $\Gamma_{\mu_1}, \Gamma_{\mu_2}$  are the free boundaries corresponding to (2.1). So, there are three cases to be considered:  $u(0) > \mu_2, \mu_1 < u(0) \leq \mu_2$  and  $u(0) \leq \mu_1$ .

We denote by  $u_\lambda^*$  the solution of (2.1) in the case  $u(0) > \mu_2$  (without a free boundary), by  $u_{\lambda, \mu_2}$  the solution in the case  $\mu_1 < u(0) \leq \mu_2$  (with one free boundary) and by  $u_{\lambda, \mu_1, \mu_2}$  the solution in the case  $u(0) \leq \mu_1$  (with two free boundaries).

**Theorem 2.1.** For  $\varepsilon \in (0, 1)$ , we have the following result:

(i) If  $\lambda < \lambda_1 := \frac{6(u_\infty - \mu_2)}{R^2}$ , then there exists a unique solution  $u_\lambda^*$  of (2.1). Moreover,

$$u_\lambda^*(0) = \frac{-\lambda}{6} R^2 + u_\infty > \mu_2,$$

i.e. the line  $(\lambda, \gamma^*(\lambda))$ , where  $u_\lambda^*(0) := \gamma^*(\lambda) := \frac{-\lambda}{6} R^2 + u_\infty$  defines a decreasing part of the bifurcation diagram (see Figure 1).

(ii) There exists  $\lambda^* > 0$  such that if  $\lambda \in [\lambda_1, \lambda^*)$ , then there exists a unique positive solution  $u_{\lambda, \mu_2}$  of (2.1) giving rise to a free boundary given by

$$r_\lambda = \frac{(\varepsilon - \frac{3}{2})}{3(\varepsilon - 1)} R \left[ 1 + 2 \cos \left( \frac{1}{3} \arccos \left( 1 + \frac{27(\varepsilon - 1)^2 \left( \frac{1}{2} R^2 - \frac{3(u_\infty - \mu_2)}{\lambda} \right)}{2(\varepsilon - \frac{3}{2})^3 R^2} \right) + \frac{4\pi}{3} \right) \right].$$

(iii) If  $\lambda \in [\lambda^*, +\infty)$ , the problem (2.1) has a unique solution  $u_{\lambda, \mu_1, \mu_2}$  with two free boundaries namely  $r_{\lambda, \mu_1}$  and  $r_{\lambda, \mu_2}$  satisfying  $0 < r_{\lambda, \mu_1} < r_{\lambda, \mu_2} < R$ .

*Proof.* First, to prove (i), we consider the case  $u(0) > \mu_2$  (without a free boundary). We have the corresponding problem

$$\begin{cases} \frac{1}{r^2} \frac{\partial}{\partial r} \left( r^2 \frac{\partial u}{\partial r} \right) = \lambda & \text{in } (0, R), \\ u(R) = u_\infty, u'(0) = 0. \end{cases}$$




Fig. 1. A qualitative description of the bifurcation curve

So,

$$u_\lambda^*(r) = \frac{\lambda}{6}(r^2 - R^2) + u_\infty, \quad r \in [0, R].$$

Since

$$u_\lambda^*(0) = \frac{-\lambda}{6}R^2 + u_\infty > \mu_2 \iff \lambda < \frac{6(u_\infty - \mu_2)}{R^2} := \lambda_1.$$

Secondly, to prove (ii), we assume that  $\mu_1 < u(0) \leq \mu_2$ . In this case, a solution take the value  $\mu_2$  at only one value of  $r$ , say  $r_\lambda$ , i.e.  $u(r_\lambda) = \mu_2$ . Hence, we have the following problem:

$$\begin{cases} \frac{1}{r^2} \frac{\partial}{\partial r} (r^2 \frac{\partial u}{\partial r}) = \lambda (\varepsilon + (1 - \varepsilon)H(u - \mu_2)) & \text{in } (0, R), \\ u(R) = u_\infty, u'(0) = 0. \end{cases} \tag{2.3}$$

In [1], we have studied the solutions of problem (2.3) for all  $\varepsilon > 0$ . Here, we reduce the study for  $\varepsilon \in (0, 1)$ . More precisely, we have the following problem in the different regions  $(0, r_\lambda)$  and  $(r_\lambda, R)$ .

$$\begin{cases} \frac{1}{r^2} \frac{\partial}{\partial r} (r^2 \frac{\partial u}{\partial r}) = \lambda \varepsilon & \text{in } (0, r_\lambda), \\ \frac{1}{r^2} \frac{\partial}{\partial r} (r^2 \frac{\partial u}{\partial r}) = \lambda & \text{in } (r_\lambda, R), \\ u(r_\lambda) = \mu_2, u'(0) = 0, u(R) = u_\infty. \end{cases}$$

Thus, the following solution is obtained:

$$u_{\lambda, \mu_2}(r) = \begin{cases} \frac{\lambda \varepsilon}{6}(r^2 - r_\lambda^2) + \mu_2, & 0 \leq r \leq r_\lambda, \\ u_\infty + \frac{\lambda}{6}(r^2 - R^2) + \frac{(r-R)r_\lambda}{(r_\lambda-R)r}(\mu_2 - u_\infty - \frac{\lambda}{6}(r_\lambda^2 - R^2)), & r_\lambda \leq r \leq R. \end{cases}$$

Using the transmission condition

$$\frac{\partial u_{\lambda, \mu_2}}{\partial r}(r_{\lambda}^-) = \frac{\partial u_{\lambda, \mu_2}}{\partial r}(r_{\lambda}^+),$$

where  $\frac{\partial u_{\lambda, \mu_2}}{\partial r}(r_{\lambda}^+)$ ,  $\frac{\partial u_{\lambda, \mu_2}}{\partial r}(r_{\lambda}^-)$  are the right and the left derivatives at the value  $r = r_{\lambda}$ , respectively. Thus,

$$\lambda = \frac{3(u_{\infty} - \mu_2)}{(\varepsilon - 1)r_{\lambda}^2 \frac{(R - r_{\lambda})}{R} + \frac{1}{2}(R^2 - r_{\lambda}^2)} := g(r_{\lambda}).$$

Then, for  $\lambda > g(0) := \lambda_1$  and  $\varepsilon \in (0, 1)$  we obtain the explicit form of  $r_{\lambda}$ :

$$r_{\lambda} = \frac{(\varepsilon - \frac{3}{2})}{3(\varepsilon - 1)} R \left[ 1 + 2 \cos \left( \frac{1}{3} \arccos \left( 1 + \frac{27(\varepsilon - 1)^2 (\frac{1}{2} R^2 - \frac{3(u_{\infty} - \mu_2)}{\lambda})}{2(\varepsilon - \frac{3}{2})^3 R^2} \right) \right) + \frac{4\pi}{3} \right].$$

A complete analysis can be found in [1].

Now, to prove (iii) we shall assume that  $u(0) \leq \mu_1$ . As in the case (ii), we look for the free boundaries  $\{(r_{\lambda, \mu_i}, \theta) : \theta \in S\}$ ,  $i = 1, 2$ , with  $u(r_{\lambda, \mu_i}) = \mu_i$ . So, we consider the corresponding problems verified by  $u$  in the different regions  $(0, r_{\lambda, \mu_1})$ ,  $(r_{\lambda, \mu_1}, r_{\lambda, \mu_2})$  and  $(r_{\lambda, \mu_2}, R)$ . On  $(0, r_{\lambda, \mu_1})$ , we have the following problem:

$$\begin{cases} \frac{1}{r^2} \frac{\partial}{\partial r} (r^2 \frac{\partial u}{\partial r}) = 0 & \text{in } (0, r_{\lambda, \mu_1}), \\ u(r_{\lambda, \mu_1}) = \mu_1, u'(0) = 0. \end{cases}$$

Then,

$$u_{\lambda, \mu_1, \mu_2}(r) = \mu_1, \quad 0 \leq r \leq r_{\lambda, \mu_1}.$$

On  $(r_{\lambda, \mu_1}, r_{\lambda, \mu_2})$ , we get the problem

$$\begin{cases} \frac{1}{r^2} \frac{\partial}{\partial r} (r^2 \frac{\partial u}{\partial r}) = \lambda \varepsilon & \text{in } (r_{\lambda, \mu_1}, r_{\lambda, \mu_2}), \\ u(r_{\lambda, \mu_2}) = \mu_2, \frac{\partial u}{\partial r}(r_{\lambda, \mu_1}^-) = \frac{\partial u}{\partial r}(r_{\lambda, \mu_1}^+) \end{cases}$$

and thus,

$$u_{\lambda, \mu_1, \mu_2}(r) = \frac{\lambda \varepsilon}{6} (r^2 - r_{\lambda, \mu_2}^2) + \frac{\lambda \varepsilon}{3} r_{\lambda, \mu_1}^3 \left( \frac{1}{r} - \frac{1}{r_{\lambda, \mu_2}} \right) + \mu_2, \quad r_{\lambda, \mu_1} \leq r \leq r_{\lambda, \mu_2}.$$

Finally, we examine the problem

$$\begin{cases} \frac{1}{r^2} \frac{\partial}{\partial r} (r^2 \frac{\partial u}{\partial r}) = \lambda & \text{in } (r_{\lambda, \mu_2}, R), \\ u(R) = u_{\infty}, \quad \frac{\partial u}{\partial r}(r_{\lambda, \mu_2}^-) = \frac{\partial u}{\partial r}(r_{\lambda, \mu_2}^+). \end{cases}$$

So,

$$\begin{aligned} u_{\lambda, \mu_1, \mu_2}(r) &= \frac{\lambda}{6} (r^2 - R^2) \\ &+ \left( \frac{\lambda(\varepsilon - 1)}{3} r_{\lambda, \mu_2}^3 - \frac{\lambda \varepsilon}{3} r_{\lambda, \mu_1}^3 \right) \left( \frac{1}{R} - \frac{1}{r} \right) + u_{\infty}, \quad r_{\lambda, \mu_2} \leq r \leq R. \end{aligned}$$

Our solutions must satisfy the conditions

$$\begin{cases} u_{\lambda, \mu_1, \mu_2}(r_{\lambda, \mu_1}^+) = \mu_1, \\ u_{\lambda, \mu_1, \mu_2}(r_{\lambda, \mu_2}^+) = \mu_2. \end{cases}$$

More precisely, we obtain the system

$$\begin{cases} \frac{\lambda \varepsilon}{6} (r_{\lambda, \mu_1}^2 - r_{\lambda, \mu_2}^2) + \frac{\lambda \varepsilon}{3} r_{\lambda, \mu_1}^3 \left( \frac{1}{r_{\lambda, \mu_1}} - \frac{1}{r_{\lambda, \mu_2}} \right) = \mu_1 - \mu_2, \\ \frac{\lambda}{6} (r_{\lambda, \mu_2}^2 - R^2) + \left( \frac{\lambda(\varepsilon-1)}{3} r_{\lambda, \mu_2}^3 - \frac{\lambda \varepsilon}{3} r_{\lambda, \mu_1}^3 \right) \left( \frac{1}{R} - \frac{1}{r_{\lambda, \mu_2}} \right) = \mu_2 - u_\infty, \end{cases} \quad (2.4)$$

where  $0 \leq r_{\lambda, \mu_1} < r_{\lambda, \mu_2} < R$ . The resolution of the system (2.4) gives the existence of the free boundaries  $r_{\lambda, \mu_1}$  and  $r_{\lambda, \mu_2}$  to conclude with the study of solutions of (2.1). The first equation in the system (2.4) can be regarded as an algebraic equation, where the unknown variable is  $r_{\lambda, \mu_2}$ . We have

$$r_{\lambda, \mu_2}^3 - \left( 3r_{\lambda, \mu_1}^2 + \frac{6(\mu_2 - \mu_1)}{\lambda \varepsilon} \right) r_{\lambda, \mu_2} + 2r_{\lambda, \mu_1}^3 = 0. \quad (2.5)$$

Taking  $r_{\lambda, \mu_1} := \eta$ , this implies that

$$\lambda = \frac{6(\mu_2 - \mu_1) r_{\lambda, \mu_2}}{\varepsilon (r_{\lambda, \mu_2}^3 - 3 \eta^2 r_{\lambda, \mu_2} + 2\eta^3)}. \quad (2.6)$$

As in [1], we introduce the auxiliary function

$$K(r) = \frac{6(\mu_2 - \mu_1) r}{\varepsilon (r^3 - 3 \eta^2 r + 2\eta^3)}, \quad r \in (\eta, R).$$

Hence,

$$K'(r) = \frac{12(\mu_2 - \mu_1)(\eta^3 - r^3)}{\varepsilon (r_{\lambda, \mu_2}^3 - 3 \eta^2 r_{\lambda, \mu_2} + 2\eta^3)^2} < 0, \quad r \in (\eta, R).$$

The function  $K$  is monotonely decreasing on  $(\eta, R)$ . Then if

$$\lambda > K(R) = \frac{6(\mu_2 - \mu_1)R}{\varepsilon (R^3 - 3\eta^2 R + 2\eta^3)} := \lambda_2,$$

the equation (2.6) has one root  $r_{\lambda, \mu_2} \in (\eta, R)$ .

So, assuming that  $\lambda > \lambda_2$  and using Cardano's method for the equation (2.5), we get

$$\begin{aligned} \Delta_2 &= (2 r_{\lambda, \mu_1}^3)^2 - \frac{4}{27} \left( 3r_{\lambda, \mu_1}^2 + \frac{6(\mu_2 - \mu_1)}{\lambda \varepsilon} \right)^3 \\ &= \frac{-8(\mu_2 - \mu_1)}{\lambda \varepsilon} \left[ 2 r_{\lambda, \mu_1}^2 \left( r_{\lambda, \mu_1}^2 + \frac{\mu_2 - \mu_1}{\lambda \varepsilon} \right) + \left( r_{\lambda, \mu_1}^2 + \frac{2(\mu_2 - \mu_1)}{\lambda \varepsilon} \right)^2 \right] < 0. \end{aligned}$$

Then, the free boundary  $r_{\lambda, \mu_2}$  is given by

$$r_{\lambda, \mu_2} = 2\sqrt{r_{\lambda, \mu_1}^2 + \frac{2(\mu_2 - \mu_1)}{\lambda\varepsilon}} \cos\left(\frac{1}{3} \arccos\left(-r_{\lambda, \mu_1}^3 \frac{1}{\sqrt{\left(r_{\lambda, \mu_1}^2 + \frac{2(\mu_2 - \mu_1)}{\lambda\varepsilon}\right)^3}}\right) + \frac{4\pi}{3}\right).$$

Now, we look for the unknown variable  $r_{\lambda, \mu_1}$  in the second equation of the system (2.4). We get

$$r_{\lambda, \mu_1} = \left(\left(\frac{3(u_\infty - \mu_2)}{\lambda\varepsilon} + \frac{r_{\lambda, \mu_2}^2 - R^2}{2\varepsilon}\right) \frac{R r_{\lambda, \mu_2}}{r_{\lambda, \mu_2} - R} + \left(\frac{\varepsilon - 1}{\varepsilon}\right) r_{\lambda, \mu_2}^3\right)^{\frac{1}{3}}.$$

Hence, the free boundary  $r_{\lambda, \mu_1}$  satisfies  $0 < r_{\lambda, \mu_1} < r_{\lambda, \mu_2}$ , if and only if the inequality  $0 < h(r) < r^2$  has a solution on  $(0, R)$ , where


$$h(r) = \left(\frac{3(u_\infty - \mu_2)}{\lambda\varepsilon} + \frac{r^2 - R^2}{2\varepsilon}\right) \frac{R}{r - R} + \left(\frac{\varepsilon - 1}{\varepsilon}\right) r^2$$

and


$$h'(r) = \frac{3(u_\infty - \mu_2)}{\lambda\varepsilon} \frac{-R}{(r - R)^2} + \frac{R}{2\varepsilon} + 2\left(\frac{\varepsilon - 1}{\varepsilon}\right) r.$$

An easy calculation shows that the function  $h'$  is decreasing on  $(0, R)$ . Then, we distinguish two cases (see Figures 2 and 3).

- (1) If  $h'(0) \leq 0$ , i.e.  $\lambda \leq \lambda_1$ , the function  $h$  is decreasing and negative on  $(0, R)$ .
- (2) If  $h'(0) > 0$ , i.e.  $\lambda > \lambda_1$ , the function  $h$  has a positive maximum in  $r^* \in (0, R)$ .  
 Moreover,  $h$  is monotone increasing on  $(0, r^*)$  and monotone decreasing on  $(r^*, R)$ .



**Fig. 2.** The graph of the function  $h$  and the square function when  $\lambda \leq \lambda_1$



**Fig. 3.** The graph of the function  $h$  and the square function when  $\lambda > \lambda_1$

So, for  $\lambda > \lambda_1$ , we get that the inequality  $0 < h(r) < r^2$  has at least one solution on  $(0, R)$ .

Finally, we assure the existence of  $\lambda^* > \max(\lambda_1, \lambda_2)$  such that when  $\lambda \in [\lambda^*, +\infty)$ , the free boundaries  $r_{\lambda, \mu_1}$  and  $r_{\lambda, \mu_2}$  satisfy  $0 < r_{\lambda, \mu_1} < r_{\lambda, \mu_2} < R$ . Thus, by continuity of the solution with respect to  $\lambda$ , there exists  $\lambda^*$  such that if  $\lambda \in [\lambda_1, \lambda^*[$ , then there exists a unique solution  $u_{\lambda, \mu_2}$  such that  $u_{\lambda, \mu_2}(r_\lambda) = \mu_2$  and when  $\lambda \in [\lambda^*, +\infty[$ , we have a unique solution with two free boundaries  $r_{\lambda, \mu_1}$  and  $r_{\lambda, \mu_2}$ . The proof of Theorem 2.1 ends.  $\square$

Let the pair  $(u, R_s)$  denote a solution of the problem (2.1)–(2.2) such that  $R_s > 0$ . Then  $R_s$  is obtained as a zero of the equation  $I(R) = 0$ , where the continuous function  $I : [0, \infty) \rightarrow \mathbb{R}$  is defined by

$$I(R) = \frac{1}{R^3} \left( \lambda \int_0^R f(u)r^2 dr - \int_0^R (\eta_1 H(u - \mu_1) + \eta_2 H(\mu_1 - u))r^2 dr \right). \quad (2.7)$$

According to Theorem 2.1, we see that we have three phases,  $u(0) > \mu_2$ ,  $\mu_1 < u(0) \leq \mu_2$  and  $u(0) \leq \mu_1$ .

The tumor remains in phase 1 (without free boundary) until  $u_\lambda^*(0) = \mu_2$  giving

$$R^* = \sqrt{\frac{6(u_\infty - \mu_2)}{\lambda}}.$$

The tumor now in phase 2 until  $u_{\lambda, \mu_2}(0) = \mu_1$  giving the radius  $R^{**}$  and the tumor moves from phase 2 into phase 3. So, when  $R < R^*$ , the tumor region is in phase 1 corresponds to the absence of free boundary (normal growth case).

When  $R^* \leq R < R^{**}$ , the tumor region is in phase 2 with one free boundary (slow growth case),  $R \geq R^{**}$  the tumor is in phase 3 corresponds to the necrotic core (with two free boundaries). Then we have the following result:

**Theorem 2.2.** *Assume that*

$$\lambda\varepsilon \geq \eta_1 \tag{2.8}$$

*is satisfied. Then there exists a unique positive value  $R_s$  such that  $I(R_s) = 0$ .*

The proof of Theorem 2.2 is based on the following lemma:

**Lemma 2.3.** *The operator  $I$  is continuously differentiable on  $(0, +\infty)$ .*

*Proof.* Consider the functions

$$\begin{cases} F_1(x_1, x_2, y) = \frac{\lambda\varepsilon}{6}(x_1^2 - x_2^2) + \frac{\lambda\varepsilon}{3} x_1^3 \left(\frac{1}{x_1} - \frac{1}{x_2}\right) - \mu_1 + \mu_2, \\ F_2(x_1, x_2, y) = \frac{\lambda}{6}(x_2^2 - y^2) + \left(\frac{\lambda(\varepsilon-1)}{3} x_2^3 - \frac{\lambda\varepsilon}{3} x_1^3\right) \left(\frac{1}{y} - \frac{1}{x_2}\right) - \mu_2 + u_\infty. \end{cases} \tag{2.9}$$

The Jacobian matrix of  $F_1, F_2$  with respect to  $x_1, x_2$  is given by

$$J = \begin{pmatrix} \lambda\varepsilon \left(x_1 - \frac{x_1^2}{x_2}\right) & \frac{\lambda\varepsilon}{3} \left(-x_2 + \frac{x_1^3}{x_2^2}\right) \\ -\lambda\varepsilon x_1^2 \left(\frac{1}{y} - \frac{1}{x_2}\right) & \frac{\lambda(3-2\varepsilon)}{3} x_2 + \lambda(\varepsilon-1) \frac{x_2^2}{y} - \frac{\lambda\varepsilon}{3} \frac{x_1^3}{x_2^2} \end{pmatrix}.$$

By (2.4), we have

$$F_1(r_{\lambda,\mu_1}, r_{\lambda,\mu_2}, R) = F_2(r_{\lambda,\mu_1}, r_{\lambda,\mu_2}, R) = 0.$$

Using the fact that  $r_{\lambda,\mu_1} < r_{\lambda,\mu_2} < R$ , we obtain that  $\det J > 0$  in the neighborhood of  $(r_{\lambda,\mu_1}, r_{\lambda,\mu_2}, R)$ . Hence, by the implicit function theorem, in the neighborhood of  $(r_{\lambda,\mu_1}, r_{\lambda,\mu_2}, R)$  there are two differentiable functions  $x_1(y)$  and  $x_2(y)$  satisfy the system (2.4). So, our free boundaries  $R \rightarrow r_{\lambda,\mu_1}(R), r_{\lambda,\mu_2}(R)$  are differentiable on  $R$ .

On the other hand, we have

$$\begin{aligned} I(R) &= \frac{1}{R^3} \left( \lambda \int_0^R (\varepsilon H(u - \mu_1) + (1 - \varepsilon) H(u - \mu_2)) r^2 dr \right. \\ &\quad \left. - \int_0^R (\eta_1 H(u - \mu_1) + \eta_2 H(\mu_1 - u)) r^2 dr \right) \\ &= \frac{1}{R^3} \left( \int_{r_{\lambda,\mu_1}}^{r_{\lambda,\mu_2}} \lambda \varepsilon r^2 dr + \int_{r_{\lambda,\mu_2}}^R \lambda r^2 dr - \int_{r_{\lambda,\mu_1}}^R \eta_1 r^2 dr - \int_0^{r_{\lambda,\mu_1}} \eta_2 r^2 dr \right) \\ &= \frac{\lambda\varepsilon(r_{\lambda,\mu_2}^3 - r_{\lambda,\mu_1}^3) + \lambda(R^3 - r_{\lambda,\mu_2}^3) - \eta_1(R^3 - r_{\lambda,\mu_1}^3) - \eta_2 r_{\lambda,\mu_1}^3}{R^3}. \end{aligned}$$

Then the operator  $I$  is differentiable on  $(0, +\infty)$ . □

**Lemma 2.4.** *Assuming that (2.8) holds, then the function  $I$  is decreasing and satisfies*

$$-\frac{\eta_2}{3} \leq I(R) \leq \frac{\lambda - \eta_1}{3} \tag{2.10}$$

*Proof.* Setting  $s = \frac{r}{R}$ , we have

$$\begin{aligned} I(R) &= \lambda \int_0^1 f(u(Rs))s^2 ds - \int_0^1 (\eta_1 H(u(Rs) - \mu_1) + \eta_2 H(\mu_1 - u(Rs)))s^2 ds \\ &= \int_0^1 g(u(Rs))s^2 ds, \end{aligned}$$

where

$$\begin{aligned} g(u) &= (\lambda\varepsilon - \eta_1)H(u - \mu_1) + \lambda(1 - \varepsilon)H(u - \mu_2) - \eta_2 H(\mu_1 - u) \\ &= \begin{cases} -\eta_2, & u < \mu_1, \\ \lambda\varepsilon - \eta_1, & \mu_1 \leq u < \mu_2, \\ \lambda - \eta_1, & u > \mu_2. \end{cases} \end{aligned}$$

Let  $U(r, R) = u(r)$ , then  $U$  is the solution of the following problem:

$$\begin{cases} \Delta_r U = \lambda f(U), & 0 < r < R, \\ U_r(0, R) = 0, & U(R, R) = u_\infty, \end{cases}$$

where

$$\Delta_r U = \frac{1}{r^2} \frac{\partial}{\partial r} \left( r^2 \frac{\partial U}{\partial r} \right).$$

Thus,  $U_R := \frac{\partial U}{\partial R}$  satisfies

$$\begin{cases} \Delta_r U_R = \frac{\partial \lambda}{\partial R} f(U), & 0 < r < R, \\ \frac{\partial}{\partial r} U_R(0, R) = 0, & U_R(R, R) = 0. \end{cases}$$

Using the fact that  $R \mapsto \lambda(R)$  is decreasing and by maximum principle, we obtain:  $R \mapsto U(r, R)$  is decreasing. The assumption (2.8) implies that the function  $g$  is increasing. Hence, we deduce that the function  $I$  is decreasing on  $(0, +\infty)$  and satisfies (2.10). This proves Lemma 2.4. □

*Proof of Theorem 2.2.* For  $R < R^*$ , we have

$$I(R) = \frac{\lambda - \eta_1}{3} > 0.$$

In [1], a simple calculation shows the existence of  $\bar{R}$  such that

$$L(\bar{R}) = \frac{1}{\bar{R}^2} \left( \lambda \int_0^{\bar{R}} \tilde{f}(u)r^2 dr - \int_0^{\bar{R}} \eta_1 r^2 dr \right) = 0,$$

where

$$\tilde{f}(u(r)) = \begin{cases} \varepsilon, & \text{if } r < r_\lambda, \\ 1, & \text{if } r \geq r_\lambda \end{cases}$$

and  $r_\lambda$  is given in Theorem 2.1. Hence,  $L(\bar{R}) = 0$  is equivalent to

$$(\lambda - \eta_1)\bar{R}^3 = -\lambda(\varepsilon - 1)r_\lambda^3 \tag{2.11}$$

So, we distinguish two cases:

- (1) if  $\bar{R} \in (0, R^{**})$ , then  $I(R) = L(R)/R$  and  $I(\bar{R}) = 0$ ,
- (2) if  $\bar{R} \in [R^{**}, +\infty)$ , then we have

$$I(\bar{R}) = \frac{1}{\bar{R}^3} \left( \frac{\lambda(\varepsilon - 1)}{3} r_{\lambda, \mu_2}^3 + \frac{\eta_1 - \lambda\varepsilon - \eta_2}{3} r_{\lambda, \mu_1}^3 + \frac{\lambda - \eta_1}{3} \bar{R}^3 \right)$$

using (2.11) and the assumption (2.8) we obtain

$$I(\bar{R}) = \frac{1}{\bar{R}^3} \left( \frac{\lambda(\varepsilon - 1)}{3} (r_{\lambda, \mu_2}^3 - r_\lambda^3) + \frac{\eta_1 - \lambda\varepsilon - \eta_2}{3} r_{\lambda, \mu_1}^3 \right) < 0.$$

So, we conclude with the existence of  $R_s$  such that  $I(R_s) = 0$  and uniqueness follows from the monotonicity of  $I$  proving in Lemma 2.4. □

### 3. TRANSIENT SOLUTION

In this section, we shall prove the global existence of a solution and the asymptotic behavior of solutions to the free boundary problem (1.4).

**Theorem 3.1.** *Assuming that (2.8) is satisfied, then for any  $R_0 > 0$  and for  $0 < \varepsilon < 1$ , the problem (1.4) has a unique global solution  $(u(r, t), R(t))$  for  $t > 0$ . Moreover,*

$$\lim_{t \rightarrow +\infty} R(t) = R_s, \quad \lim_{t \rightarrow +\infty} u(r, t) = u_{\lambda, \mu_1, \mu_2},$$

where  $(u_{\lambda, \mu_1, \mu_2}, R_s)$  is the stationary solution of the problem (2.1)–(2.2).

To prove theorem 3.1, we use the following lemma:

**Lemma 3.2** ([25]). *Let consider the problem*

$$\begin{cases} x'(t) = x(t)f(x(t)), & t > 0, \\ x(0) = x_0 > 0. \end{cases}$$

*Assuming that  $f$  is continuously and differentiable decreasing function on  $(0, +\infty)$ . If there exists a unique positive constant  $x_s$  such that  $f(x_s) = 0$ , then*

$$\lim_{t \rightarrow +\infty} x(t) = x_s.$$

*Proof of Theorem 3.1.* We know from Section 2 that problem (1.4) admits a unique solution  $u(r, R(t))$ . So, for  $t > 0$ , we determine  $R(t)$  by solving the following problem:

$$\begin{cases} R'(t) = R(t)I(R(t)), & t > 0, \\ R(0) = R_0, \end{cases} \tag{3.1}$$

where  $I$  is given by (2.7). Using Lemmas 2.3 and 2.4, then the problem (3.1) has a unique global solution satisfying

$$R_0 e^{\left(\frac{-\eta_2}{3}\right)t} \leq R(t) \leq R_0 e^{\left(\frac{\lambda-\eta_1}{3}\right)t}, \quad t > 0.$$

Now, because  $I(R)$  is decreasing function and from theorem 2.2, there exists  $R_s$  such that the equation  $I(R_s) = 0$ . Then, by Lemma 3.2, we get

$$\lim_{t \rightarrow +\infty} R(t) = R_s.$$

So,  $\lim_{t \rightarrow +\infty} u(r, t) = u_{\lambda, \mu_1, \mu_2}$ . This concludes the proof of Theorem 3.1. □

#### 4. PERTURBATION RESULT

This section is devoted to study the problem (1.3) when  $\Omega(t)$  verifies (C). Let  $r_1, r_2$  denote the values  $r_{\lambda, \mu_1}$  and  $r_{\lambda, \mu_2}$  of theorem 2.1. We look for the free boundaries on the form  $r_1 + b_1(\theta)$  and  $r_2 + b_2(\theta)$ , where  $b_1, b_2$  are the perturbations caused by  $\beta$ . Hence, we consider the following problem:

$$\begin{cases} \Delta u = \lambda \left( \varepsilon H(u - \mu_1) + (1 - \varepsilon) H(u - \mu_2) \right) & \text{in } \Omega_\beta, \\ u = \bar{u}_\infty & \text{on } \partial\Omega_\beta, \end{cases} \tag{4.1}$$

where  $\Omega_\beta = B(0, R + \beta(\theta)) \subset \mathbb{R}^3$  and  $\bar{u}_\infty$  is close to  $u_\infty$ .

As in [1], we define the set of admissible surfaces in  $\Omega_\beta$  by

$$S_\beta = \{f \in C(S) : (f(\theta), \theta) \in \Omega_\beta \text{ for } \theta \in S\}.$$

For  $i = 1, 2$ , we consider the functions  $\psi_i \in S_\beta$  and the sets

$$\omega_{\psi_i} = \{(r, \theta) \in \Omega_\beta : r < \psi_i(\theta)\}, \quad i = 1, 2.$$

In the following proposition, we formulate nonlinear equations for  $\psi_1, \psi_2$  and prove that by solving them, we can solve the problem (4.1).

**Proposition 4.1.** *For  $\varepsilon \in (0, 1)$ , there exists  $\lambda^* > 0$  such that if  $\lambda \geq \lambda^*$ , the following problem*

$$\begin{cases} \Delta u = \lambda \left( \varepsilon \chi_{\Omega_\beta \setminus \omega_{\psi_1}} + (1 - \varepsilon) \chi_{\Omega_\beta \setminus \omega_{\psi_2}} \right) & \text{in } \Omega_\beta, \\ u = \bar{u}_\infty & \text{on } \partial\Omega_\beta \end{cases} \tag{4.2}$$

has a unique solution  $u_\beta \in C^{1, \alpha}(\overline{\Omega_\beta})$  with  $\alpha = 1 - \frac{3}{p}$ ,  $p > 3$ . Moreover, if  $u_\beta(\psi_i(\theta), \theta) = \mu_i$  for  $i = 1, 2$  and  $\bar{u}_\infty > \mu_2 > \mu_1$ , then  $u_\beta$  is a solution of (4.1).

*Proof.* We have  $\chi_{\Omega_\beta \setminus \omega_{\psi_i}} \in L^p(\Omega_\beta)$ ,  $p > 1$ ,  $i = 1, 2$ . From [19], there exists a unique solution  $u_\beta$  of (4.2) in  $W^{2,p}(\Omega_\beta)$ . For  $p > 3$ ,  $W^{2,p}(\Omega_\beta) \subset C^{1,\alpha}(\overline{\Omega_\beta}, \mathbb{R})$  with  $\alpha = 1 - \frac{3}{p}$ .

If we prove the existence of functions  $\psi_i$  such that  $u_\beta(\psi_i(\theta), \theta) = \mu_i$  for  $i = 1, 2$ , then  $u_\beta$  will be a solution of the following problems:

$$\begin{cases} \Delta u_\beta = 0 & \text{in } \omega_{\psi_1}, \\ u_\beta = \mu_1 & \text{on } \partial\omega_{\psi_1}, \end{cases} \quad \begin{cases} \Delta u_\beta = \lambda\varepsilon & \text{in } \omega_{\psi_2} \setminus \omega_{\psi_1}, \\ u_\beta = \mu_1 & \text{on } \partial\omega_{\psi_1}, \\ u_\beta = \mu_2 & \text{on } \partial\omega_{\psi_2}, \end{cases} \quad \begin{cases} \Delta u_\beta = \lambda & \text{in } \Omega_\beta \setminus \omega_{\psi_2}, \\ u_\beta = \mu_2 & \text{on } \partial\omega_{\psi_2}, \\ u_\beta = \bar{u}_\infty & \text{on } \partial\Omega_\beta. \end{cases}$$

Using the fact that  $\bar{u}_\infty > \mu_2 > \mu_1$  and the maximum principle, we obtain that  $u_\beta$  satisfies

$$\begin{cases} \Delta u_\beta = \lambda(\varepsilon H(u_\beta - \mu_1) + (1 - \varepsilon)H(u_\beta - \mu_2)) & \text{in } \Omega_\beta, \\ u_\beta = \bar{u}_\infty & \text{on } \partial\Omega_\beta. \end{cases} \quad \square$$

As in [1], the variation of the domain  $\Omega_\beta$  suggests the use of an appropriate transformation that maps the changing domain  $\Omega_\beta$  to a constant domain:

$$\begin{aligned} T_\beta: \Omega_\beta &\rightarrow \Omega_0 = B(0, R), \\ (r, \theta) &\mapsto (\bar{r}, \theta) = \left(r + r \frac{\beta}{R}, \theta\right), \end{aligned}$$

where  $(r, \theta)$  is the coordinates in  $\Omega_\beta$  and  $(\bar{r}, \theta)$  the coordinates in  $\Omega_0$ .

For a small  $\beta$ , the transformation  $T_\beta$  is a diffeomorphism of class  $C^2$  of the domain  $\Omega_\beta$  into  $\Omega_0$  and transforms  $S_\beta$  into  $S_0$ , hence

$$T_\beta(\psi_i(\theta), \theta) = (f_i(\theta), \theta),$$

where  $f_i \in S_0$ . So, the mapping  $T_\beta$  transforms the problem (4.2) and the equation  $u_\beta(\psi_i(\theta), \theta) = \mu_i$  for  $i = 1, 2$  to the problem

$$\begin{cases} \Delta \bar{u} + \delta_\beta \bar{u} = \lambda(\varepsilon \chi_{\Omega_0 \setminus \omega_{f_1}} + (1 - \varepsilon) \chi_{\Omega_0 \setminus \omega_{f_2}}) & \text{in } \Omega_0, \\ \bar{u} = \bar{u}_\infty & \text{on } \partial\Omega_0 \end{cases} \quad (4.3)$$

and the equation

$$\bar{u}(f_i(\theta), \theta) = \mu_i, \quad i = 1, 2. \quad (4.4)$$

where

$$\begin{aligned} \delta_\beta &= \frac{\beta}{R} \left(2 + \frac{\beta}{R}\right) \frac{\partial^2}{\partial \bar{r}^2} \\ &+ \frac{2\beta}{\bar{r}R} \frac{\partial}{\partial \bar{r}} \frac{1}{\bar{r}^2} \left[ a_{ij}(\theta) \left[ \frac{r}{R} \frac{\partial \beta}{\partial \theta_j} \left( \frac{\partial^2}{\partial \bar{r} \partial \theta_i} + \frac{1}{R(1 + \frac{\beta}{R})} \frac{\partial \beta}{\partial \theta_i} \frac{\partial}{\partial \bar{r}} + \frac{r}{R} \frac{\partial \beta}{\partial \theta_i} \frac{\partial^2}{\partial \bar{r}^2} \right) \right. \right. \\ &\quad \left. \left. + \frac{r}{R} \frac{\partial^2 \beta}{\partial \theta_j \partial \theta_i} \frac{\partial}{\partial \bar{r}} + \frac{r}{R} \frac{\partial \beta}{\partial \theta_i} \frac{\partial^2}{\partial \theta_j \partial \bar{r}} \right] + b_i(\theta) \left[ \frac{r}{R} \frac{\partial \beta}{\partial \theta_i} \frac{\partial}{\partial \bar{r}} \right] \right] \end{aligned}$$

for  $a_{i,j}, b_i \in C^2(S)$ ,  $i, j = 1, 2$ , and  $f_i \in S_0$ .

Thus, to solve the problem (4.1) it is sufficient to prove the existence of  $f_1, f_2$  satisfying (4.4). We can proceed as follows.

First, we give the existence of  $f_2(\theta)$  for  $\theta \in S$ . We denote by  $\bar{u}_2$  the restriction of the solution  $\bar{u}$  in the set  $\Omega_0 \setminus \omega_{f_2}$ . So, we have

$$\begin{cases} \Delta \bar{u}_2 + \delta_\beta \bar{u}_2 = \lambda(\varepsilon + (1 - \varepsilon) \chi_{\Omega_0 \setminus \omega_{f_2}}) & \text{in } \Omega_0, \\ \bar{u}_2 = \bar{u}_\infty & \text{on } \partial\Omega_0. \end{cases} \tag{4.5}$$

The same technique used in [1] can be applied and the implicit function theorem gives the existence of  $f_2(\theta), \theta \in S$  such that

$$\bar{u}_2(f_2(\theta), \theta) = \mu_2.$$

Hence, from the construction and uniqueness of the solution we conclude that

$$\bar{u}(f_2(\theta), \theta) = \mu_2.$$

Now, consider the following problem

$$\begin{cases} \Delta \bar{u}_1 + \delta_\beta \bar{u}_1 = \lambda\varepsilon \chi_{\Omega_0 \setminus \omega_{f_1}} & \text{in } \Omega_0, \\ \bar{u}_1 = \bar{u}_\infty & \text{on } \partial\Omega_0. \end{cases}$$

We define the operator  $J : \mathbb{R}^+ \times S_0 \times \mathbb{R}^+ \times D \rightarrow C(S, \mathbb{R})$  by

$$J(\bar{u}_\infty, f_1, \mu_1, \beta) = \bar{u}_1(f_1(\theta), \theta) - \mu_1,$$

where  $D$  is the neighborhood of zero in  $C(S)$ .

So, we obtain

$$\begin{aligned} J(\bar{u}_\infty, f_1, \mu_1, \beta) &= \bar{u}_\infty \int_S P(f_1(\theta), \theta, \theta') d\theta' + \lambda\varepsilon \int_S \int_{f_1(\theta')}^R G(f_1(\theta), \theta, r', \theta') r'^2 dr' d\theta' \\ &\quad - \int_{\Omega_0} \delta_\beta \bar{u}_1(r', \theta') G(f_1(\theta), \theta, r', \theta') dr' d\theta' - \mu_1, \end{aligned}$$

where  $P$  is the Poisson kernel and  $G$  is the Green function for the Laplacian in  $\Omega_0$ .

By the same argument as in [1] and using the implicit function theorem in the neighborhood of  $(\bar{u}_\infty, r_1, \mu_1, 0)$ , we have the existence of a function  $f_1$  depending on  $\bar{u}_\infty, \mu_1$ , where  $\beta$  satisfies

$$J(\bar{u}_\infty, f_1(\bar{u}_\infty, \mu_1, \beta), \mu_1, \beta) = 0.$$

So, we conclude that

$$\bar{u}(f_1(\theta), \theta) = \mu_1.$$

Moreover, in the region  $\omega_{f_1}$ , we have the following result:

**Lemma 4.2.** *Let  $\gamma$  be a Jordan curve in  $\{x \in \Omega_0 : \bar{u}(x) = \mu_1\}$ , where  $\omega$  is the interior of  $\gamma$ . Then*

$$\omega \subset \{x \in \Omega_0 : \bar{u}(x) = \mu_1\}.$$

*Proof.* Suppose that there exists  $x_0 \in \omega$  such that  $\bar{u}(x_0) < \mu_1$ . Let

$$\omega_1 = \{x \in \Omega_0 : \bar{u}(x) < \mu_1\}$$

which is an open set (by continuity of  $\bar{u}$ ). So,  $\bar{u}$  verifies

$$\begin{cases} \Delta u = 0 & \text{in } \omega_1, \\ u = \mu_1 & \text{on } \partial\omega_1. \end{cases}$$

By the maximum principle we get  $\bar{u} \geq \mu_1$  which is a contradiction.

So, let  $u$  be a solution of the problem (2.1), then the set  $\{x \in \Omega : u(x) = \mu_1\}$  is a ball of radius  $r_1 \in (0, R)$ .

Moreover, by the above perturbation method, we conclude that  $w_{f_1}$  is a perturbation of a ball of radius  $r_1 + f_1(\theta)$ ,  $\theta \in S$ . □

### 5. FINAL COMMENTS

In this paper, we have studied the existence of stationary solutions for problem (1.3) when the domain is a ball. After, reducing the free boundary problem (1.3) into an initial problem of ordinary differential equation, we study the asymptotic behavior of solutions. Finally, using a rigorous perturbative approach, we derive a characterization of the free boundary under perturbation of the boundary condition and a smooth boundary of the domain. However, some open questions related to our problem still remain. Here, we give some interesting problems:

(1) A more general form of the problem (2.1) can be regarded as

$$\begin{cases} \Delta u = \lambda \sum_{i=1}^n \phi_i H(u - \mu_i) & \text{in } \Omega(t), \\ u = \bar{u}_\infty & \text{on } \partial\Omega(t), \end{cases} \tag{5.1}$$

where  $\phi_i$  are positive constants for  $i = 1, \dots, n$ . To the best of our knowledge, no investigation has been devoted to such problem in the literature. Recently, in [3] the second author studies a similar problem of (5.1) and shows that under suitable conditions there exists a solution with a convex level set. We refer to [3] for other results. Moreover, the problem (5.1) can be reformulated as an equivalent free boundary problem. For  $t > 0$ , we have

$$\begin{cases} \Delta u = \lambda \sum_{i=1}^n \phi_i \chi_{\Omega(t) \setminus \omega_{\mu_i}(t)} & \text{in } \Omega(t), \\ u = \bar{u}_\infty & \text{on } \partial\Omega(t), \end{cases} \tag{5.2}$$

where  $\chi_\omega$  denotes the characteristic function of the set  $\omega$ . There is a mathematical challenge in the study of problem (5.2) and some technical details will be presented elsewhere.

- (2) Certainly, for other models, we conjecture that the results of our work is still true for  $n$  free boundaries (the nonlinearity  $f$  can be written as  $f(u) = \sum_{i=1}^n \phi_i H(u - \mu_i)$ ).
- (3) The obtained bifurcation diagram can be very important to study the question of the stability of solution of problem (1.1). We recall that the stability results obtained by the Crandall–Rabinowitz theorem [10] can not be applied in our work. This question requires more attention.

### Acknowledgements

*The authors are thankful to the honorable reviewers for their valuable suggestions and comments, which improved the paper.*

### REFERENCES

- [1] A. Abdelouahab, S. Bensid, *Perturbation analysis in a free boundary problem arising in tumor growth model*, arXiv:2303.15828.
- [2] R. Araujo, D.L.S. McElwain, *A history of the study of solid tumour growth: The contribution of mathematical modelling*, Bull. Math. Biol. **66** (2004), 1039–1091.
- [3] S. Bensid, *A free boundary problem for a discontinuous semilinear elliptic equations in convex ring*, Complex Var. Elliptic Equ. (2024), 1–10.
- [4] S. Bensid, J.I. Díaz, *Stability results for discontinuous nonlinear elliptic and parabolic problems with a S-shaped bifurcation branch of stationary solutions*, Discrete Contin. Dyn. Syst. Ser. S **22** (2017), 1757–1778.
- [5] S. Bensid, J.I. Díaz, *On the exact number of monotone solutions of a simplified Budyko climate model and their different stability*, Discrete Contin. Dyn. Syst. Ser. S **24** (2019), 1033–1047.
- [6] A. Borisovich, A. Friedman, *Symmetry-breaking bifurcations for free boundary problem*, Indiana Univ Math. J. **54** (2005), 927–947.
- [7] H.M. Byrne, *Mathematical Biomedicine and Modeling Avascular Tumor Growth*, Mathematics and Life Sciences, De Gruyter, **1** (2012), 277–303.
- [8] H.M. Byrne, M.A.J. Chaplain, *Growth of nonnecrotic tumors in the presence and absence of inhibitors*, Math. Biosci. **131** (1995), 151–181.
- [9] H.M. Byrne, M.A.J. Chaplain, *Growth of necrotic tumors in the presence and absence of inhibitors*, Math. Biosci. **135** (1996), 187–216.
- [10] M.G. Crandall, P.H. Rabinowitz, *Bifurcation from simple eigenvalues*, J. Funct. Anal. **8** (1971), 321–340.
- [11] S. Cui, *Analysis of a free boundary problem modeling tumor growth*, Acta. Math. Sin. (Engl. Ser.) **21** (2005), 1071–1082.
- [12] S. Cui, A. Friedman, *Analysis of a mathematical model of the growth of necrotic tumors*, J. Math. Anal. Appl. **255** (2001), 636–677.
- [13] E. DiBenedetto, *Partial Differential Equations*, Springer Science, Business Media New York, 1995.

- [14] M. Fontelos, A. Friedman, *Symmetry-Breaking bifurcations for free boundary problems in three dimensions*, *Asymptot. Anal.* **35** (2003), 187–206.
- [15] A. Friedman, B. Hu, *Asymptotic stability for a free boundary problem arising in tumor models*, *J. Differ. Equ.* **227** (2006), 598–639.
- [16] A. Friedman, B. Hu, *Bifurcation for stability to instability for a free boundary problem arising in a tumor model*, *Arch. Rational Mech. Anal.* **180** (2006), 293–330.
- [17] A. Friedman, K. Lam, *Analysis of a free boundary tumor model with angiogenesis*, *J. Differ. Equ.* **259** (2015), 7636–7661.
- [18] A. Friedman, F. Reitich, *Analysis of a mathematical model for the growth of tumors*, *J. Math. Biol.* **38** (1999), 262–284.
- [19] D. Gilbarg, N. Trudinger, *Elliptic Partial Differential Equations of Second Order*, Springer-Verlag, New York, 1983.
- [20] H.P. Greenspan, *Models for the growth of a solid tumor by diffusion*, *Stud. Appl. Math.* **51** (1972), 317–340.
- [21] H.P. Greenspan, *On the growth and stability of cell cultures and solid tumors*, *J. Theor. Biol.* **56** (1976), 229–242.
- [22] Y. Huang, Z. Zhang, B. Hu, *Bifurcation from stability to instability for a free boundary tumor model with angiogenesis*, *Discret Contin. Dyn. Sys.* **39** (2019), 2473–2510.
- [23] D.L.S. McElwain, L.E. Morris, *Apoptosis as a volume loss mechanism in mathematical models of solid tumor growth*, *Math. Biosci.* **39** (1978), 147–157.
- [24] C. Müller, *Spherical Harmonics*, Lecture Notes in Math., vol. 17, Springer, Berlin, Heidelberg, New York, 1966.
- [25] S. Xu, *Stability of solutions to a free boundary problem for tumor growth*, *Int. J. Differ. Equ.* **4** (2014), Article ID 427547.
- [26] J. Wu, *Stationary solutions of a free boundary problem modeling the growth of vascular tumors with Gibbs–Thomson relation*, *J. Differ. Equ.* **260** (2016), 5875–5893.
- [27] J. Wu, F. Zhou, *Asymptotic behavior of solutions of a free boundary problem modeling the growth of tumors with fluid-like tissue under the action of inhibitors*, *Trans. Amer. Math. Soc.* **365** (2013), 4181–4207.

Ahlem Abdelouahab  
ahlemabdelouahab98@gmail.com

University of Tlemcen  
Faculty of Sciences  
Department of Mathematics  
Dynamical Systems and Applications Laboratory  
B.P. 119, Tlemcen 13000, Algeria

Sabri Bensid (corresponding author)  
edp\_sabri@yahoo.fr

University of Tlemcen  
Faculty of Sciences  
Department of Mathematics  
Dynamical Systems and Applications Laboratory  
B.P. 119, Tlemcen 13000, Algeria

*Received: December 21, 2023.*

*Revised: April 20, 2024.*

*Accepted: May 10, 2024.*